## NAT Breakfast Task

# Name:

Set-up:



#### Evaluation:

Left		Right	
toast (minimum 6 slices)		toaster	
jam with loosened screw cap		butter/margarine	
cup		1 knife	
teabags in a carton		1 table spoon	
		1 tea spoon	
		kettle filled with water	
		sugar in container	
	/4		/7

### Lateralized attention score = contralesional – ipsilesional proportion:

Notes:

#### Accomplishment Score:

Toast:
--------

<ul> <li>toasted bread (1)</li> <li>apply butter <i>or</i> jam to the bread (1)</li> <li>apply butter <i>and</i> jam to the bread (2)</li> </ul>	/3
Tea: <ul> <li>switch on the kettle (1)</li> <li>put the teabag in the cup (1)</li> </ul>	
□ infuse water (1) □ add sugar (1)	/4
Sum:	/7

#### Date:

Error Score:	Number:
<ul> <li>toasts more than one slice of bread:</li> </ul>	
<ul> <li>mixes up the order:</li> </ul>	
$\Box$ fill in water then plug in the kettle	
$\Box$ infuse water, then turn on the kettle	
$\Box$ turn on the toaster without bread inside	
$\square$ butter the bread first and then put it in the toaster	
$\square$ apply jam to bread and then butter it	
$\Box$ omit to stir the tea	
$\square$ omit to spread the butter and/or jam	
<ul> <li>use one ingredient extremely excessive or sparsely:</li> </ul>	
<ul> <li>typical errors of apraxia:</li> </ul>	
$\Box$ spread with the finger	
$\Box$ spread with the spoon	
$\Box$ put the bread on top of the toaster	
$\Box$ press the bread with the hand in the toaster	
$\Box$ take butter with the spoon (for jam ok!)	
$\Box$ take sugar with the knife	
$\Box$ empty the sugar directly in the cup	
□ put jam in the cup	
$\Box$ put sugar on the bread	
$\Box$ put the teabag on the bread	

Error Score = Sum of errors:

/19

#### NAT Score:

Accomplishment Score	Error Score	Naturalistic Action Task Score	Interpretation
7	0-1	6	
7	> 1	5	no apraxia
6	0-1	4	
6	> 1	3	mild apraxia
4–5	0-1	2	moderate apraxia
4–5	> 1	1	severe apraxia
0–3	≥0	0	